

Research Article

Evaluation of Aphrodisiac Activity of *Turnera aphrodisiaca*

Suresh Kumar^{1*}, Reecha Madaan¹ and Anupam Sharma²

¹S.D. College of Pharmacy, Barnala-148 101, Punjab, India

²University Institute of Pharmaceutical Sciences, Panjab University, Chandigarh

ABSTRACT

Turnera aphrodisiaca Ward (family Turneraceae) is reputed as an aphrodisiac throughout the world. Despite a long tradition of use as an aphrodisiac, no systematic work has ever been carried out on this plant to validate its traditional use. In the present investigation, various extracts (petroleum ether, chloroform, methanol and water), alkaloidal fraction, and volatile oil of *T. aphrodisiaca* were screened for aphrodisiac activity in mice. Mounting behaviour was taken as a parameter to screen aphrodisiac activity. Chloroform extract exhibited significant activity at a dose of 200 mg/kg, p.o. while methanol extract showed aphrodisiac activity at a lower dose, i.e., 50 mg/kg, p.o.. Volatile oil of *T. aphrodisiaca* was found to be devoid of aphrodisiac activity. Qualitative phytochemical screening showed the presence of alkaloids in chloroform and methanol extracts. Therefore, the alkaloidal fraction was isolated from aerial parts of *T. aphrodisiaca*, and tested for aphrodisiac activity at dose levels of 25, 50, 75, or 100 mg/kg, p.o. A dose dependent increase in activity was observed upon acute as well as subacute administration of alkaloidal fraction.

Keywords: Anti-anxiety; Aphrodisiac; Turneraceae; *Turnera aphrodisiaca*

INTRODUCTION

Turnera aphrodisiaca Ward (synonym *T. diffusa* Willd., family Turneraceae) is commonly known as 'Damiana'. The leaves of *T. aphrodisiaca* have been used traditionally as a stimulant, aphrodisiac, tonic, diuretic, nerve tonic, laxative, and in kidney, menstrual, and pregnancy disorders⁹. The British Herbal Pharmacopoeia (1983)⁵ lists specific indications for Damiana as anxiety neurosis associated with impotency, and includes other indications as depression, nervous dyspepsia, atonic constipation and coital inadequacy. Damiana has achieved some repute in the treatment of sexual impotence where it is used in conjunction with strychnine, phosphorus or some other stimulant¹³. The leaf infusion of Damiana has been used in the diseases related to the gastrointestinal and respiratory system⁶, reproductive organs¹⁶, and for the treatment of gonorrhoea¹⁰. Mother tincture of Damiana is an important homoeopathic medicine for the treatment of sexual debility, and nervous prostration⁴. *T. aphrodisiaca* has been reported to contain cyanogenic glycoside tetraphyllin B¹⁸; flavonoid gonzalitosin I⁷; arbutin³; damianin¹⁹; tricosan-2-one, hexacosanol⁸; volatile oil containing α -pinene, β -pinene, p-cymene, and 1,8-cineole³; and β -sitosterol⁷.

Aqueous extract of *T. aphrodisiaca* whole plant has been reported to exhibit significant hypoglycaemic activity in alloxan-diabetic male mice¹⁵. In another set of experiments, Aguilara et al. (1998)¹ reported that a decoction of *T. aphrodisiaca* leaves possesses significant hypoglycaemic activity in rabbits upon oral administration. Aqueous extract of the plant has also been reported to exhibit sexual stimulating activity in sexually sluggish male rats at a dose of 1 ml/kg².

Authors have reported that amongst various extracts (petroleum ether, chloroform, methanol and water) of *T. aphrodisiaca* aerial parts, only methanol extract (25 mg/kg, p.o.) exhibited significant anti-anxiety activity on elevated plus maze apparatus¹¹. An anxiolytic constituent apigenin has been isolated from methanol extract of *T. aphrodisiaca* aerial parts using bioactivity-guided fractionation¹². Despite a long tradition of use of *T. aphrodisiaca* as aphrodisiac, only preliminary work was carried out by Arletti et al. on this plant. Therefore, various extracts viz., petroleum ether (60°-80°C), chloroform, methanol and water, alkaloidal fraction, and volatile oil of *T. aphrodisiaca* were evaluated for aphrodisiac activity in mice.

Materials and Methods

Plant Material

T. aphrodisiaca Ward aerial parts were procured from Rati Ram Nursery, Village Khurrampur, district Saharanpur (U.P.) in the month of August 2002. Identity of the plant was confirmed by the Head, NISCAIR, New Delhi.

Animals

Male and non-oestrous female Laca mice (either sex) were bred at the Central Animal House, Panjab University, Chandigarh. The animals were allowed a standard pellet diet and water *ad libitum*. Groups of five mice (20 - 24 g) were used in all sets of experiments. The animals were fasted for 18 h before use.

Solvents

Petroleum ether (60° - 80°C), chloroform (Ranbaxy Laboratory Chemicals) and methanol (S.D. Fine Chemicals Pvt.), all of LR grade, distilled under normal atmospheric pressure were employed for extraction of the plant material.

Recovery of solvents

Solvents from extracts were recovered under reduced pressure using Buchi 461 Rotary vacuum evaporator

*Corresponding author: Prof Suresh Kumar

S.D. College of Pharmacy Barnala-148 101, Punjab, India

Ph. No.: 01679-243505

Fax : 01679-243505

E-mail: thakur_pu@yahoo.com



Fig: *Turnera aphrodisiaca* Ward and were preserved in a vacuum desiccator containing anhydrous silica gel blue.

Extract	Yield (% w/w)
Petroleum ether	3.61
Chloroform	2.19
Methanol	12.10
Water	5.50

Mounting behaviour

To observe the libido-oriented mounting behaviour, non-oestrous female mice were paired with treated male mice²⁰. The male mouse assuming the copulatory position over the female was considered as a mount. After the administration of *T. aphrodisiaca* extracts, alkaloidal fraction, or volatile oil, the behaviour of the animals was observed for 3 h (9.00 a.m. - 12.00 p.m.). Males were placed individually after p.o. (i.p. in case of volatile oil) administration of *T. aphrodisiaca* in a clear aquarium (9.00 a.m.) and were allowed to

acclimatize for 15 min. After that, a non-oestrous female was introduced into the arena. The number of mounts were recorded for 15 min (9.15 a.m. - 9.30 a.m.). Females were separated for next 60 min interval and were reintroduced again for 15 min, between 10.30 a.m. - 10.45 a.m., during which mounts were recorded. Similarly, the third interaction was done between 11.45 a.m. - 12.00 p.m., and mounts were recorded.

Vehicle

Distilled water + Tween 80 (5%) was used as vehicle for preparing the suspension of various test doses of different extracts. Test doses of volatile oil were prepared using olive oil as vehicle.

Statistics

The results have been expressed as mean \pm standard deviation (S.D.). The test doses were compared with control by analysis of variance (ANOVA) followed by Fischer's LSD test¹⁷.

Preparation of extracts

Powdered aerial parts (250 g) of *T. aphrodisiaca* were successively extracted in Soxhlet apparatus using the solvents in order of increasing polarity viz., petroleum ether (60° - 80°C), chloroform, and methanol. The marc was air dried, and water extract was obtained by boiling with distilled water for 2 h. The aqueous extract was filtered, concentrated and dried in an oven at 40° - 50°C. Table 1 shows the yield of various extracts. Every extract was then subjected to biological evaluation for aphrodisiac activity in mice at various dose levels, i.e., 50, 100, 200, or 400 mg/kg, p.o. Mounting behaviour was taken as a parameter to evaluate aphrodisiac activity. Volatile oil was obtained from aerial parts (200 g) of the plant by water distillation using clevenger apparatus, and evaluated for aphrodisiac activity at the doses of 50, 100, 200, or 400 mg/kg, i.p. Since

Table 2:Aphrodisiac activity of various extracts of *T. aphrodisiaca* aerial parts

Treatment	Dose (mg/kg, p.o.)	Average ⁿ no. of mounts			
		9:15-9:30 a.m.	10:30-10:45 a.m.	11:45 a.m.-12:00 p.m.	a.m.-12:00 p.m.
Control [Distilled water + Tween 80 (5%)]	Vehicle	2.2 \pm 0.45	3.0 \pm 0.71	0.8 \pm 0.45	
Petroleum ether extract	50	2.0 \pm 0.71	2.6 \pm 0.55	0.6 \pm 0.55	
	100	1.8 \pm 0.45	3.0 \pm 0.71	0.8 \pm 0.84	
	200	1.8 \pm 0.84	1.6 \pm 0.89	1.0 \pm 0.71	
	400	2.0 \pm 0.71	2.2 \pm 0.84	1.0 \pm 0.71	
Chloroform extract	50	2.2 \pm 0.45	4.4 \pm 0.89	2.2 \pm 0.84	
	100	3.8 \pm 0.84	7.0 \pm 1.00*	2.4 \pm 0.55	
	200	4.6 \pm 0.89*	9.2 \pm 1.64*	4.2 \pm 1.10*	
	400	4.4 \pm 1.14*	8.8 \pm 1.30*	4.4 \pm 1.14*	
Methanol extract	50	3.2 \pm 0.84	9.4 \pm 1.34*	3.8 \pm 0.84*	
	100	3.0 \pm 0.71	1.6 \pm 0.89	0.0 \pm 0.00	
	200	3.2 \pm 0.84	0.0 \pm 0.00	0.0 \pm 0.00	
	400	2.4 \pm 0.55	0.0 \pm 0.00	0.0 \pm 0.00	
Water extract	50	2.6 \pm 0.55	2.8 \pm 0.84	0.8 \pm 0.45	
	100	2.0 \pm 0.71	2.2 \pm 0.45	1.0 \pm 0.71	
	200	2.0 \pm 0.71	2.8 \pm 1.64	1.4 \pm 0.39	
	400	1.8 \pm 0.45	2.0 \pm 0.71	1.2 \pm 0.45	

N = 5; *P<0.05 vs. control; ANOVA followed by Fischer's LSD test.

qualitative phytochemical screening showed the presence of alkaloids in chloroform and methanol extracts. Therefore, the alkaloidal fraction was also evaluated for aphrodisiac activity. The alkaloidal fraction was isolated from aerial parts of *T. aphrodisiaca* by following method: Aerial parts (2 kg) of *T. aphrodisiaca* were treated with lime, and then Soxhlet extracted with chloroform. The chloroform extract was concentrated to 1/4th of its original volume under reduced pressure. It was then partitioned in a separator using 5 × 50 ml of 2% acidulated water (HCl - water). The aqueous fraction was basified using NaOH solution to pH 8 – 9 followed by partitioning with chloroform (5 × 50 ml). The chloroform fraction was rich in alkaloids. This alkaloidal fraction was dried, and subjected to aphrodisiac activity upon acute administration of 25, 50, 75, or 100 mg/kg, p.o. Various doses, i.e., 25, 50, 75, or 100 mg/kg of alkaloidal fraction were also administered twice daily (9:00 a.m. and 5:00 p.m.) for nine days in mice. The aphrodisiac activity was observed on the 10th day of administration of respective dose.

Results

Tables 2-4 show the average number of mounts after acute administration (single dose) of various extracts viz., petroleum ether, chloroform, methanol and water (50, 100, 200, or 400 mg/kg, p.o.), volatile oil (50, 100, 200, or 400 mg/kg, i.p.) and alkaloidal fraction (25, 50, 75, or 100 mg/kg, p.o.) respectively. Table 5 shows the average number of mounts after subacute administration of alkaloidal fraction (25, 50, 75, or 100 mg/kg, p.o.).

Discussion

As is evident from table 2, chloroform and methanol extracts were found to be active. Chloroform extract exhibited maximum aphrodisiac activity at the dose of 200 mg/kg while methanol extract at the dose of 50 mg/kg during 2nd h of test with respect to control. The aphrodisiac activity declined with the passage of time, i.e., during 3rd h of the test as the bioactive constituent(s) seem to have undergone a metabolic degradation. Mice did not show any activity at higher doses of methanol due to prevalence of sedative effects. Volatile oil did not exhibit aphrodisiac

Table 4: Aphrodisiac activity of alkaloidal fraction of *T. aphrodisiaca* upon acute administration.

Treatment	Dose (mg/kg, p.o.)	Average ⁿ no. of mounts		
		9:15-9:30 a.m.	10:30-10:45 a.m.	11:45 a.m.-12:00 p.m.
Control [Distilled water + Tween 80 (5%)]	Vehicle	2.2 ± 0.84	2.8 ± 0.89	1.4 ± 0.89
Alkaloidal fraction	25	4.0 ± 0.71	6.0 ± 1.22*	2.2 ± 0.71
	50	4.2 ± 1.10	9.4 ± 1.14*	2.4 ± 0.55
	75	4.0 ± 1.22	10.0 ± 1.87*	3.2 ± 0.84
	100	4.4 ± 1.14	11.2 ± 1.92*	4.0 ± 0.89*

n = 5; *P<0.05 vs. control; ANOVA followed by Fischer's LSD test.

Table 5: Aphrodisiac activity of alkaloidal fraction of *T. aphrodisiaca* upon subacute (9 days) administration.

Treatment	Dose (mg/kg, p.o.)	Average ⁿ no. of mounts on 10 th day		
		9:15-9:30 a.m.	10:30- 10:45 a.m.	11:45 a.m.-12:00 p.m.
Control [Distilled water + Tween 80 (5%)]	Vehicle	2.6 ± 0.84	3.8 ± 1.10	1.4 ± 0.89
Alkaloidal fraction	25	4.6 ± 1.67*	12.8 ± 2.59*	4.0 ± 0.89*
	50	5.2 ± 1.09*	14.8 ± 3.03*	3.8 ± 0.84*
	75	5.8 ± 0.84*	14.4 ± 2.30*	5.2 ± 0.71*
	100	6.0 ± 1.22*	15.0 ± 2.55*	5.6 ± 2.07*

n = 5; *P<0.05 vs. control; ANOVA followed by Fischer's LSD test.

Table 3: Aphrodisiac activity of volatile oil of *T. aphrodisiaca* aerial parts.

Treatment	Dose (mg/kg, i.p.)	Average ⁿ no. of mounts		
		9:15-9:30 a.m.	10:30-10:45 a.m.	11:45 a.m.-12:00 p.m.
Control [Olive oil]	Vehicle	2.4 ± 0.55	3.2 ± 0.84	1.2 ± 0.45
Volatile oil	50	2.2 ± 0.45	2.0 ± 0.71	1.0 ± 0.71
	100	2.0 ± 1.22	1.6 ± 0.55	1.0 ± 0.71
	200	2.2 ± 1.30	2.6 ± 0.55	1.4 ± 0.89
	400	1.6 ± 0.55	1.8 ± 0.84	0.8 ± 0.45

n = 5; *P<0.05 vs. control; ANOVA followed by Fischer's LSD test.

activity at any of the doses tested (Table 3). A dose dependent increase in aphrodisiac activity was observed in alkaloidal fraction upon acute as well as subacute administration during 2nd h of interaction (Table 4-5). Male mice showed maximum aphrodisiac activity upon sub-acute administration of alkaloidal fraction. This observation infers that *T. aphrodisiaca* improves reproductive organs upon long term use. One more interesting observation is that crude methanol extract and purified alkaloidal fraction exhibited significant aphrodisiac activity at similar dose, i.e., 50 mg/kg. This observation suggests that one or more constituent(s) in combination with bioactive alkaloid possess synergistic beneficial effects on reproductive organs. Bioactivity directed fractionation, using various chromatography techniques, work is in progress to isolate bioactive alkaloid from *T. aphrodisiaca*.

Acknowledgements

Authors duly acknowledge the financial assistance provided by University Grants Commission, New Delhi to Suresh Kumar for this research work.

References

1. Aguilar FJA, Ramos RR, Gutierrez SP, Contretras AA, Weber CCC, Saenz JLF (1998): Study of the anti-hyperglycaemic effect of plants used as antidiabetics. *J Ethnopharmacol* 61: 101-110.
2. Arletti R, Benelli A, Cavazzuti E, Scarpetta G, Bertolini A (1999): Stimulating property of *Turnera diffusa* and *Pfaffia paniculata* extracts on the sexual behavior of male rats. *Psychopharmacology* 143: 15-19.
3. Auterhoff H, Hackle HP (1968): Components of damiana drug. *Archiv Pharm* 301: 537-544.
4. Boericke W (1988). *Pocket Manual of Homoeopathic Materia Medica*. New Delhi, India, B. Jain Publisher Private Limited, p. 659.
5. British Herbal Pharmacopoeia (1983). West Yorks, British Herbal Medicine Association, p. 29.
6. Caceres A (1996): *Turnera aphrodisiaca*. In: Giron L, Caceres A, eds., *Plantas de Uso Medicinal en Guatemala*. Editorial Universitaria San Carlos de Guatemala, pp. 160-162.
7. Dominguez XA, Hinojosa M (1976): Mexican medicinal plants. XXVIII. Isolation of 5-hydroxy-7, 3', 4'-trimethoxy flavone from *Turnera diffusa*. *Planta Med* 30: 68-71.
8. Fryer FA (1965): Chemical investigation of damiana (*Turnera diffusa*). *Specialities* 1: 21-23.
9. Hocking GM, Thomas CC (1955). *A Dictionary of Terms in Pharmacognosy*. Illinois, USA, Springfield, p. 234.
10. Koch L (1936): Drug collection from Bolivia systematically, anatomically and chemically examined. *Arch Pharmacol* 274: 343-369.
11. Kumar S, Sharma A (2005): Anti-anxiety activity studies of various extracts of *Turnera aphrodisiaca* Ward. *J Herb Pharmacother* 5: 13-21.
12. Kumar S, Sharma A (2006): Apigenin – The anxiolytic constituent of *Turnera aphrodisiaca* Ward. *Pharm Biol* 44: 84-90.
13. Osol A, Farrar GF, Leuallen EE, Youngken HW, Detweiler DK (1947). *Dispensary of United States of America* (24th edition). Philadelphia, J. B. Lippincott Company, pp. 1422-1423.
14. Parfitt K (1999). *Martindale, the Complete Drug Reference* (32nd edition). 1 Lambeth High Street, London, Pharmaceutical Press, p. 1570.
15. Perez RM, Ocegueda A, Munoz JL, Avita JG, Morrow WW (1984): A study of the hypoglycaemic effect of some Mexican plants. *J Ethnopharmacol* 12: 253-262.
16. Saggese D (1959). *Medicinal Herbs of Argentina* (10th edition). Rosario, Argentina, Antoghazzi & Co., pp. 1-189.
17. Scheffer WC (1980). *Statistics for the Biological Sciences*. Philippines, Addison-Wesley Publishing Company, Inc., pp. 121-141.
18. Spencer KC, Seigler DS (1981): Tetraphyllin B from *Turnera diffusa*. *Planta Med* 43: 175-178.
19. Steinmetz EF (1960): *Damianae folia*. *Acta Phytotherapeut* 7: 1-2.
20. Subramoniam A, Madhavachandran R, Rajasekharan S, Pushpangadan P (1997): Aphrodisiac property of *Trichopus zeylanicus* extract in male mice. *J Ethnopharmacol* 57: 21-27.